

Map User Notes: Hazelnuts Suitability Map (Using Climate Futures Projections)

Summary:

The mapping aims to classify land suitable for growing Hazelnuts according to seven suitability classes: “1.0 Well suited”, “1.1 Well suited (with soil management)”, “2.0 Suitable”, “2.1 Suitable (with soil management)”, “3.0 Moderately suitable”, “3.1 Moderately suitable (with soil management)”, “3.2 Moderately suitable (with frost/heat management)”, “3.3 Moderately suitable (with soil, frost/heat management)”, and “4.0 Unsuitable”. These are produced from a set of pre-determined rules in accordance to a suitability matrix (refer to page 3) developed from reviewing existing literature in conjunction with industry consultation and ground-truthing. They are based on soil and climatic influences on Hazelnut production using data garnered from soil cores and temperature logger records collected from hundreds of site locations across the state. The mapping assumes water for crop irrigation is available and therefore not a limiting factor to production.

[Climate Futures](#) projections were incorporated into the suitability mapping framework to assess projected changes to frost/heat risk severity, chill hours, growing degree days and mean temperature/rainfall for years [2030](#) and [2050](#) (refer to [metadata](#) for more information). This compliments the current [Hazelnuts Suitability Map](#) developed as part of the [‘Water for Profit’](#) program which was based on recent historical climate observations (Refer [here](#) for more information).

How is suitability determined?

The overall suitability rating is determined using a most-limiting-factor approach, where the lowest rated parameter becomes the overall suitability rating. As an example, a location may possess the following characteristics:

<u>Soil/Climate parameter</u>	<u>Threshold</u>	<u>Rating</u>
Soil depth:	>60cm	(Well suited)
pH:	6-7	(Well suited)
Electrical conductivity:	<0.15	(Well suited)
Texture (%Clay)	10-30%	(Well suited)
Drainage:	Well drained	(Well suited)
Stones:	<10%	(Well suited)
Slope:	10-20%	(Moderately suitable)
Risk of having no extreme temperatures in winter	>4/5years	(Well suited)
Chill hours:	>1500	(Well suited)
Frost risk (after bud-break):	<1/5years	(Well suited)
Minimum temperature for fertilisation:	>21°C	(Well suited)
Growing degree days (GDD):	<700	(Unsuitable)
Heat stress in January:	<30°C	(Well suited)
Extreme heat risk (December to January):	<1/5years	(Well suited)
Rainfall:	<50mm	(Well suited)
Overall rating:		4.0 Unsuitable

In this case, the overall suitability is classified as “4.0 Unsuitable” due to GDD not meeting the required threshold of at least 700 GDD units therefore limiting suitability due to this classification. Alternatively, if for example, GDD was ‘>800 (Well suited)’, the overall rating would then revert to “3.0 Moderately suitable”, due to the Slope parameter ‘10-20% (Moderately suitable)’ now limiting the rating to this classification (refer to suitability matrix on page 3 for

rule-set). A “1.0 Well suited” rating is achieved if all attributes are rated ‘Well suited’. Note that designated conservation/protection areas as well as existing urbanised/residential zones and major waterbodies were automatically classified as “4.0 Unsuitable”.

Improving suitability (with management)

In some circumstances, suitability ratings can be improved if measures are applied to mitigate against certain ‘soft’ limitations/constraints. Addressing these constraints (also known as ‘Manageable constraints’) using practises such as, for example, applying agricultural lime to counteract acidic soils, can result in a parcel of land becoming more suitable for a given suitability classification. As an example, a location may possess the following characteristics:

<u>Soil/Climate parameter</u>	<u>Threshold</u>	<u>Rating</u>
Soil depth:	>60cm	(Well suited)
pH:	5.2-5.5*	(Moderately suitable)
Electrical conductivity:	<0.15	(Well suited)
Texture (%Clay)	10-30%	(Well suited)
Drainage:	Well drained	(Well suited)
Stones:	<10%	(Well suited)
Slope:	<10%	(Well suited)
Risk of having no extreme temperatures in winter	>4/5years	(Well suited)
Chill hours:	>1500	(Well suited)
Frost risk (after bud-break):	<1/5years	(Well suited)
Minimum temperature for fertilisation:	>21°C	(Well suited)
Growing degree days (GDD):	>800	(Well suited)
Heat stress in January:	<30°C	(Well suited)
Extreme heat risk (December to January):	<1/5years	(Well suited)
Rainfall:	<50mm	(Well suited)
Overall rating		2.1 Suitable (with soil management)

In normal circumstances, the above rating would be determined as “3.0 Moderately suitable” due to pH limiting suitability to this classification. However, in this case, the overall suitability is reclassified to “2.1 Suitable (with soil management)” because the low pH rating, on most occasions, can be readily mitigated (e.g. by applying agricultural lime to neutralise soil acidity) therefore rendering the land more suitable for production. In this regard, the following ‘Manageable constraints’ can be actively managed/mitigated to allow improved Hazelnut growing conditions.

- Soil depth – Installation of raised soil beds can improve soil depth prior to planting.
- pH – Undertaking an active liming regime over several seasons can effectively reduce soil acidity over time. (refer: <http://dpiwwe.tas.gov.au/agriculture/land-management-and-soils/soil-management/soil-ph-liming>)
- Drainage – Installation of surface and sub-surface drainage lines can carry excess water off agricultural land and prevent water logging. (refer: <http://dpiwwe.tas.gov.au/agriculture/land-management-and-soils/soil-management/land-drainage>)
- Stones– Stone removal via machine and/or manual picking prior to planting can effectively improve the soil for crop production.

Each manageable constraint (as mentioned above) is denoted by the symbol * in the suitability matrix (page 3) to indicate where management of soil and/or climate impediments ought to be considered to improve growing conditions and meet the corresponding suitability rating. Note that ‘Manageable constraints’ can be identified in the attribute table of the map layer itself on LISTmap (refer to example on page 5 to identify how constraints are recognised in LISTmap).

Supplementary information for managing soils:

- Soil management website (DPIPWE):

<http://dipwe.tas.gov.au/agriculture/land-management-and-soils/soil-management>

- Information for soil management for Tasmanian farmers:
<http://dipwe.tas.gov.au/Documents/Soil-Guide.pdf>
- Information for managing Tasmania's cropping soils:
<http://dipwe.tas.gov.au/Documents/Managing-Tasmanias-Cropping-Soils-2000.pdf>

Supplementary information for growing hazelnuts:

- Hazelnut Growing Factsheet:
http://dipwe.tas.gov.au/Documents/WfW-Hazelnut_factsheet.pdf

Suitability Rule Matrix - Hazelnuts

Suitability rating	Soil Depth	pH (top 15cm)	Electrical conductivity dS/m	Soil texture (top 15cm - % clay)	Soil drainage	Stone abundance (>200mm diameter, top 15cm)	Slope (of land, % rise)	Extreme air temperature in winter (Chance of having no days <-7°C in June to August) (using Climate Futures Projections)	Chill hours (0-7°C, May – August) (using Climate Futures Projections)	Frost risk after bud break (Chance of having at least one day where Tmin <-3°C in October) (using Climate Futures Projections)	Air temperature at fertilisation (December mean maximum air temperature) (using Climate Futures Projections)	Growing Degree Days (GDD) (above a base of 10°C, 1 November to 30 April) (using Climate Futures Projections)	Heat stress (January mean maximum air temperature) (using Climate Futures Projections)	Extreme heat risk (Chance of having at least 2 days where Tmax >35°C in December to January) (using Climate Futures Projections)	Rainfall at harvest (Monthly mean for March) (using Climate Futures Projections)
1.0 Well suited	>60cm	6-7	<0.15	10–30%	Well drained, Moderately well drained	<10%	<10%	Occurs >4/5 years	>1500	Occurs <1/5 years	>21°C	>800	<30°C	Occurs <1/5 years	<50mm
1.1 Well suited (with soil management)	50-60cm*	5.5-6*	<0.15	10–30%	Imperfectly drained*	10-20%*	<10%	Occurs >4/5 years	>1500	Occurs <1/5 years	>21°C	>800	<30°C	Occurs <1/5 years	<50mm
2.0 Suitable	50-60cm	5.5-6	<0.15	30-50%	Imperfectly drained	10-20%	<10%	Occurs 3/5 to 4/5	1000-1500	Occurs <1/5 years	>20°C	>800	<30°C	Occurs <1/5 years	<50mm
2.1 Suitable (with soil management)	30-50 cm*	5.2–5.5*	<0.15	30-50%	Imperfectly drained	10-20%	<10%	Occurs 3/5 to 4/5	1000-1500	Occurs <1/5 years	>20°C	>800	<30°C	Occurs <1/5 years	<50mm
3.0 Moderately suitable	30-50 cm	5.2–5.5	<0.15	30-50%	Imperfectly drained	10-20%	10-20%	Occurs 2/5 to 3/5	800-1000	Occurs 1/5 – 1/3 years	>19°C	700-800	<30°C	Occurs 1/5 – 1/3 years	50-75mm
3.1 Moderately suitable (with soil management)	20-30 cm*	5.2-5.5	<0.15	30-50%	Imperfectly drained	20-50%*	10-20%	Occurs 2/5 to 3/5	800-1000	Occurs 1/5 – 1/3 years	>19°C	700-800	<30°C	Occurs 1/5 – 1/3 years	50-75mm
4.0 Unsuitable	<20cm	<5.2	>0.15	>50% or <10%	Poor, Very poorly drained	>50%	>20%	Occurs <2/5 years	<800	Occurs >1/3 years	<19°C	<700	>30°C	Occurs >1/3 years	>75mm

Manageable constraints are denoted by the symbol *

Definitions of Suitability Classes (in relation to the suitability matrix)

1.0 Well suited: Land having no significant soil or climatic limitations to sustained Hazelnut production where risk of significant crop loss due to adverse conditions are unlikely.

1.1 Well suited (with soil management): Land having no significant soil or climatic limitations to sustained Hazelnut production where risk of significant crop loss due to adverse conditions are unlikely. However, minor soil impediments with regard to soil depth, pH, drainage and/or stone abundance ought to be managed if land is to be well suited for sustained production.

2.0 Suitable: Land having only minor soil or climatic limitations that will not significantly reduce productivity where any risk of crop loss is possible, but inherently low.

2.1 Suitable (with soil management): Land having only minor soil or climatic limitations that will not significantly reduce productivity where any risk of crop loss is possible, but inherently low. However, soil impediments with regard to soil depth, pH, and/or stone abundance must be managed if land is to be suitable for sustained production. Failure to do so may result in an increased risk of crop loss.

3.0 Moderately suitable: Land having soil or climatic limitations that is likely to impact on sustained productivity where risk of significant crop loss is possible in some years.

3.1 Moderately suitable (with soil management): Land having soil or climatic limitations that is likely to impact on sustained productivity where risk of significant crop loss is possible in some years. Major soil impediments with regard to soil depth, pH and/or drainage must be managed. Failure to do so may result in significant crop loss and would be considered marginal or even unsuitable for large-scale production if constraints were left unattended.

4.0 Unsuitable: Land having soil and climatic limitations that are too severe for sustained Hazelnut production and will so reduce benefits, or increase required inputs, that this expenditure may not justify. Risk of crop loss is high.

LISTmap instructions

The Hazelnut suitability map is a digital layer that can be manually interrogated within [LISTmap](#). Any location can be enquired within Tasmania to provide location specific parameters pertaining to Hazelnut suitability. To interrogate a location, simply click on any location whilst the Hazelnuts suitability layer is active in the table of contents panel and a window will appear listing important attributes. An example is given below:

The screenshot shows the LISTmap interface with a map of Tasmania. A red pin is placed on the map, and an 'Identify Results' window is open. The window displays the following information:

One feature found in one layer
Enterprise Suitability - Hazelnuts (one feature)

Feature		
User Notes	You should read the User Notes via the hyperlink to the right prior to using this information.	User Notes (.pdf Document)
Overall Suitability Rating	3.0 Moderately suitable	
Constraints	Chill hours, Rainfall	
Soil Depth (cm)	>60cm (Well suited)	
pH (top 15cm)	5.5-6 (Suitable)	
Electrical conductivity (ECse) dS/m	<0.15dS/m (Well suited)	
Soil Texture (Clay %)	30-50% (Suitable)	
Soil Drainage	Imperfect (Suitable)	

POI: GDA94 MGA55 : 472440E, 5428995N

Identify Options

The legend on the right shows the following suitability ratings:

- 1.1 Well suited (with soil management)
- 2.0 Suitable
- 2.1 Suitable (with soil management)
- 3.0 Moderately suitable
- 3.1 Moderately suitable (with soil management)
- 4.0 Unsuitable

The window provides information such as the “Overall Suitability Rating” as well as other useful information including:

- ‘Constraints’ - provides a list of attributes that limits a classification to a particular suitability rating;
- ‘Manageable Constraints’ - provides a lists of soft limitation attributes than should be managed in order to meet the suitability rating (refer to page 1 & 2).

In addition, information pertaining to each soil and climate attribute (according to the classifications provided in the suitability matrix on page 3) are also provided and appear below the overall suitability rating and constraint fields (as shown in the example above). This allows users to ‘drill down’ and view site-specific information regarding soil and climate requirements that relate to Hazelnut production for any point of interest in Tasmania.

Vulnerable soils

Listed within the suitability layer (below the individual soil/climate fields) are ratings pertaining to soil vulnerability hazards including:

- Sodicity (ratings of high, moderate, low, nil);
- Salinity (ratings of high, moderate, low, nil);
- Water Erosion (ratings of extreme, very high, high, moderate, low, very low, nil);
- Water Logging (ratings of extreme, very high, high, moderate, low, very low, nil); and
- Wind Erosion (ratings of high, moderate, low, nil).

Information in regards to managing vulnerable soils can be obtained here:

https://nrmdatalibrary.dpipwe.tas.gov.au/FactSheets/WfW/ListMapUserNotes/Vulnerable_soils.pdf

Note that vulnerable soils categories are complementary material to the mapping and does not contribute to the “Overall Suitability Rating”.

Flood risk

Also listed within the suitability layer are categories relating to risk of flood inundation, as provided by flood plain layer within the LIST Hydrography dataset

(http://listdata.thelist.tas.gov.au/public/LIST_Hydrographic_Information.pdf).

Ratings of 'High', 'Moderate' or 'Nil' are listed that correspond to 'Severe', 'Moderate' or 'Nil' flooding potential.

Note that flood risk categories are complementary material to the mapping and does not contribute to the "Overall Suitability Rating".

Additional Information

For information about using LISTmap, please consult the help document:

<http://listdata.thelist.tas.gov.au/public/outgoing/sif/listmaphelp.pdf>

For technical information relating to the dataset, please consult the [metadata](#).

For information relating to irrigation investment, please consult the Investing in Irrigation web-portal:

<http://dppwe.tas.gov.au/agriculture/investing-in-irrigation>

References

Baldwin, B 2015, 'The growth and productivity of Hazelnut cultivars', PhD thesis, University of Sydney, Sydney NSW.

(<https://ses.library.usyd.edu.au/handle/2123/12892>)

Contact

Natural Assets Spatial Intelligence Section

171 Westbury Road

PROSPECT TAS 7250

Phone: 03 6777 2220

Email: nasis@dppwe.tas.gov.au

Caution and Disclaimer

The information and material in LISTmap (including the enterprise suitability map layer for Hazelnuts and accompanying soil and climate input datasets, i.e. "material") is based on computer modelling of the potential suitability of Hazelnuts to a given area and, as such, there are inherent uncertainties in the results. While every effort has been made to ensure the material is accurate, the Crown in Right of Tasmania ("Crown") provide no warranty, guarantee or representation that the material is accurate, complete, up to date, non-infringing or fit for a particular purpose. All suitability assessments are based upon the assumption that water for crop irrigation is available and therefore is not a limiting factor. Furthermore, the Crown expressly disclaim all and any legal liability and responsibility whatsoever arising from or connected with: (a) the accuracy, reliability, validity, currency or completeness of the material; (b) the consequences of anything done or omitted to be done by any person, either in whole or in part, in reliance of the material. The material does not take into account personal circumstances. The material is made available on the understanding that the Crown are not providing professional advice and that users of this material should undertake site-specific investigations and research and obtain appropriate professional advice relevant to their particular circumstances. The relevant maps that form part of the material have been prepared at 1:50,000 scale (landscape level). These maps consider only soil and climate constraints and do not take into account other parameters or any legislative, regulatory and/or policy requirements of Federal, State or Local Governments that apply to the land in question and/or which could affect the proposed land use or agricultural enterprise.